

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁴ : A61K 9/64, C07K 3/20, 15/14 C07K 17/00, 17/02, G01N 33/567 G01N 33/574, 33/577</p>	<p>A1</p>	<p>(11) International Publication Number: WO 89/11273</p> <p>(43) International Publication Date: 30 November 1989 (30.11.89)</p>
<p>(21) International Application Number: PCT/US89/02240</p> <p>(22) International Filing Date: 22 May 1989 (22.05.89)</p> <p>(30) Priority data: 196,986 20 May 1988 (20.05.88) US</p> <p>(71) Applicant: LA JOLLA CANCER RESEARCH FOUNDATION [US/US]; 10901 North Torrey Pines Road, La Jolla, CA 92037 (US).</p> <p>(72) Inventors: RUOSLAHTI, Erkki, I. ; ENGVALL, Eva ; P.O. Box 1054, Rancho Santa Fe, CA 92067 (US). GEHLSSEN, Kurt, R. ; 7646-16 Palmilla Drive, San Diego, CA 92122 (US).</p> <p>(74) Agents: CAMPBELL, Cathryn et al.; Pretty, Schroeder, Brueggemann & Clark, 444 South Flower Street, Suite 2000, Los Angeles, CA 90071 (US).</p>		<p>(81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: ADHESION RECEPTOR FOR LAMININ AND ITS USE</p> <p>(57) Abstract</p> <p>An adhesion receptor for laminin is provided. The receptor is isolated from cell or tissue extracts and fractionated on an affinity column composed of cell attachment-promoting fragments of laminin coupled to Sepharose in the presence of divalent cations. This receptor can be used to prepare specific antibodies for the analysis of the amount of laminin receptor expressed by cells and has other applications in cellular and tumor biology.</p>		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	ML	Mali
AU	Australia	FR	France	MR	Mauritania
BB	Barbados	GA	Gabon	MW	Malawi
BE	Belgium	GB	United Kingdom	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IT	Italy	RO	Romania
BJ	Benin	JP	Japan	SD	Sudan
BR	Brazil	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic			SN	Senegal
CG	Congo	KR	Republic of Korea	SU	Soviet Union
CH	Switzerland	LI	Liechtenstein	TD	Chad
CM	Cameroon	LK	Sri Lanka	TG	Togo
DE	Germany, Federal Republic of	LU	Luxembourg	US	United States of America
DK	Denmark	MC	Monaco		
ES	Spain	MG	Madagascar		

ADHESION RECEPTOR FOR LAMININ AND ITS USEFIELD OF INVENTION

This invention relates generally to the fields of cell adhesion systems and specifically to an adhesion receptor 5 for laminin.

BACKGROUND OF THE INVENTION

Much of what goes on at the cell surface relates to the recognition by the cell of substances around it. A prime example is the binding of soluble hormones by the cell and 10 its response to such binding. Another important aspect of cell surface recognition is the interaction of the cell with insoluble structures that surround it. Such a structure can be the surface of another cell or the extracellular matrix.

15 Although the interactions of cells with one another and with extracellular matrices are not well understood, they play an important role in the life of the cell. For example, cell-matrix and cell-cell interactions seem to tell a cell where in the body it is supposed to be or where 20 it is supposed to go if it has to migrate. A particularly intriguing example is nerve cells which send out processes to appropriate positions and thereby form connections to distant parts of the body. Positional signalling is obviously impaired in cancer since tumors invade and 25 disseminate into parts of the body inappropriate for their cell of origin. In fact, as our understanding of the cell-matrix and cell-cell interactions increases and their exploration becomes possible, this field is very likely to develop into a new, important frontier of medicine.

30 Both proteins and carbohydrates at the cell surface can participate in the cell-matrix and cell-cell interactions. Extracellular matrices are composed of an insoluble

meshwork of protein and carbohydrate that is laid down by cells and fills most of the intercellular spaces. Matrices in different locations in the body consist of different combinations of collagens, proteoglycans, elastin, 5 hyaluronic acid and various glycoproteins such as fibronectin and laminin. Virtually all of the extracellular matrix glycoproteins and collagens that have been identified interact with cells.

The most readily observable result of the interaction of 10 cells with the extracellular matrix molecules is cell adhesion. The adhesive properties of the extracellular matrix proteins can be easily demonstrated in vitro by plating cells onto a surface coated with extracellular matrix material or with one of the purified matrix 15 proteins. The cells will rapidly adhere to such a surface and spread on it. However, the adhesive proteins not only promote adhesion, they also stimulate cell migration. When confronted with limiting concentrations of an adhesive protein applied as a gradient on a surface, cells move 20 toward the higher concentration.

More complex ways in which extracellular matrices influence cells are to promote cell differentiation, survival and growth. One of the extracellular matrix proteins, laminin, has particularly striking effects on 25 cells. This protein, which is present in the specialized extracellular matrix sheets, basement membranes, promotes the attachment and migration of cells and plays a role in differentiation and tumor metastasis. Laminin also promotes and guides the growth of nerve cell processes, or 30 neurites.

These interactions between cells and laminin are thought to be mediated by cell surface receptors which function as the adhesion receptor for laminin. The full nature of the receptor, or receptors, that mediate the 35 effects of laminin on cells is still unknown.

There thus exists a need to identify and isolate laminin adhesion receptors. The availability of isolated receptor will allow production of antibodies specific for this receptor, which could be used, for example, to assay for the expression of laminin receptors at the surface of tumor cells. Compounds, such as recombinant protein fragments that bind to the receptor, can be used to reproduce the activity of laminin or to inhibit the adhesion of cells to laminin-containing structures. Moreover, there exists a need to be able to target liposomes to specific tissues for therapeutic or other purposes. The present invention satisfies these needs and provides additional advantages as well.

SUMMARY OF THE INVENTION

The present invention provides a substantially pure mammalian laminin adhesion receptor characterized in that it comprises two subunits, α and β , and interacts with laminin and cell attachment-promoting fragments of laminin from the portion of laminin that consists of the COOH-terminal portions of its polypeptides. The interaction with laminin or cell-attachment promoting fragments of laminin is divalent-cation dependent. The larger (α) subunit is reactive with anti- α_3 antisera.

In one aspect, the invention provides a method of isolating and purifying the laminin adhesion receptor. In another aspect of the present invention, monoclonal and polyclonal antibodies are prepared against the isolated receptor and their reactivity is assayed against the purified receptor. Because no other receptors are known to contain α_3 subunit, antibodies against the larger alpha subunit are specific to the laminin receptor, whereas some antibodies against the smaller beta subunit can react with the fibronectin receptor and other related cell adhesion receptors. The anti-alpha subunit antibodies are useful in determining the amount of laminin receptor which is expressed by a given cell type. Further selection by cell

attachment assays provides anti-alpha subunit antibodies that can inhibit the attachment of cells to laminin. Such antibodies prevent the invasion of tumor cells through amniotic membrane tissue in an in vitro assay.

5 In another aspect of the invention, the receptor is used to provide compounds that can reproduce or inhibit the function of the laminin receptor. In a further aspect of the present invention, the cell surface receptors are incorporated into the membranes of liposomes. Such
10 liposomes are then used to target the contents of the liposomes to tissues containing laminin.

Brief Description of the Drawings

Figure 1. Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) analysis of fractions eluted
15 from the laminin affinity matrix. RuGli glioblastoma cells were surface-labeled and extracted, and the extract fractionated as described in Example I on a Sepharose column containing covalently bound, cell-attachment-promoting fragments of laminin. Aliquots of each fraction
20 were analyzed by SDS-PAGE under nonreducing conditions, using autoradiography for visualization of protein bands. Lanes 1-13 show fractions from the EDTA elution of the column. The molecular weight markers were: myosin, 200 kD; beta-galactosidase, 116 kD; phosphorylase B, 94 kD;
25 bovine serum albumin, 67 kD; ovalbumin, 43 kD.

Figure 2. SDS-PAGE analysis of the RuGli laminin receptor (lane 1) and the fibronectin receptor (lane 2) under nonreducing (NR) and reducing (R) conditions. The fibronectin receptor was obtained from the same cell
30 extract as the laminin receptor by using fibronectin cell attachment fragment-Sepharose as the affinity matrix using the method of Pytela et al. Cell 40:548, (1985), which is incorporated herein by reference. The conditions of the electrophoresis were the same as in Figure 1. Arrows
35 indicate the light chains of the alpha subunits.

Figure 3. Immunoblot analysis of laminin receptor. The fibronectin (A) and laminin (B) receptors isolated from RuGli cells were electrophoresed under nonreducing conditions and transferred to nitrocellulose filters, and the filters were incubated with rabbit anti-fibronectin receptor antibodies (Argraves et al. J. Cell Biol. 105:1183, 1987, which is incorporated herein by reference) followed by detection of the bound antibodies with goat anti-rabbit IgG conjugated to horseradish peroxidase.

10 Figure 4. Immunoblot analysis of laminin and fibronectin receptors. The fibronectin (Lanes 1, 3) and laminin (Lanes 2, 4) receptors isolated from RuGli cell extracts were electrophoresed under nonreducing conditions and transferred to nitrocellulose filters. The filters
15 were incubated with either rabbit antiserum against the fibronectin receptor absorbed with Sepharose-coupled laminin receptor (Lanes 1, 2) or with a rabbit anti-laminin receptor antiserum absorbed with Sepharose-coupled fibronectin receptor (Lanes 3, 4). Bound antibodies were
20 detected with goat anti-rabbit IgG conjugated to horseradish peroxidase. The anti-fibronectin receptor antibody used for absorption has been described (Pytela et al., Meth. Enzymol., 144:475-489 (1987), which is incorporated herein by reference), the anti-laminin
25 receptor antiserum was prepared by immunizing with receptor purified from RuGli cells as described in Example I.

Figure 5. Binding of laminin receptor-containing liposomes to substrates. Microtiter wells were coated with various proteins and unoccupied binding sites on the
30 plastic were blocked by incubating the wells with bovine serum albumin. The binding or inhibition of binding of receptor liposomes to the wells was then tested as described in Example II. The results of six experiments are expressed as a percentage of maximal binding. The mean
35 plus standard error after subtraction of background from BSA attachment is shown: LM, human laminin; FN, fibronectin; IV, type IV collagen; LM + LM fragments, LM +

20 μ g of chymotryptic laminin fragments; LM + RGD, LM + 1mg/ml GRGDSP peptide; LM + YIGSR, LM + 1mg/ml YIGSR peptide; fibronectin receptor liposomes (open bars); LMR, laminin receptor liposomes (shaded bars).

5 Figure 6. Radioreceptor binding assay. Purified iodinated laminin receptors isolated from various human (MG-63, A431) and rat (RuGli, NRK-49F) cells were allowed to incubate with various proteins coated onto microtiter wells. The wells were washed and bound receptor was
10 solubilized in 1% SDS in TBS and quantitated in a gamma counter. The results are expressed as a percentage of receptor bound, with laminin binding equal to 100 percent. The bars represent receptors from different cell types as indicated in the Figure-LM, human laminin; FN, human
15 fibronectin; CIV, Type IV collagen; CI, Type I collagen; BSA, bovine serum albumin.

Figure 7. Inhibition of laminin receptor binding in the radio-receptor binding assay. Data represent the percentage of laminin receptor bound to laminin-coated
20 microtiter wells in the presence of various inhibitors compared to uninhibited laminin binding which was taken as 100 percent. Laminin receptors from A-431 cells (sparsely striped bars) and the RuGli cells (densely striped bars) were used. The inhibitors tested were laminin and
25 fibronectin fragments and unlabeled receptor (described under Example VIII). The RuGli receptor was not tested with two of the inhibitors (NA).

Detailed Description of the Invention

The present invention relates to a mammalian adhesion
30 receptor for laminin. Laminin is known to have a role in both the differentiation of normal cells and the metastasis of malignant ones. The isolation of this receptor and antibodies reactive with it can be used to detect the presence of the laminin receptor on the surface of tumor

cells. The ability of cells to bind to laminin has been found to correlate with their invasive and metastatic capacity.

To isolate and purify the mammalian laminin adhesion receptor, extracts of surface-labeled mammalian cells were fractionated on immobilized laminin fragments that are capable of promoting cell attachment. Preferably such cells are malignant cells of neural derivation, although other mammalian cell types can be used as well. For example, rat glioblastoma cells (RuGli) were first surface-iodinated with lactoperoxidase and dissolved in octylglucoside. The RuGli cells attach to and spread on a plastic surface coated laminin. Addition of Mn^{2+} to the cell suspension during attachment assay enhances the attachment of the cells to the laminin-coated surface. By applying the same purification method, a receptor can be isolated from other cell types and tissues, such as human osteosarcoma MG-63 cells or placental tissues.

Laminin fragments maintaining cell-attachment promoting activity were isolated generally by the method of Engvall et al., J. Cell Biol. 103:2457 (1986), which is incorporated herein by reference. Various modifications can be made to this method, including the use of intact laminin or fragments generated with proteases other than pepsin and chymotrypsin. An affinity matrix was prepared by coupling the laminin fragments to cyanogen-bromide (CNBr)-activated Sepharose (Sigma, St. Louis, MO). Specific elution was effected by treating the column with EDTA (ethylene diaminetetraacetic acid). All buffers, except the elution buffer, contained 1mM Mn^{2+} to facilitate the binding of the receptor to the affinity matrix. The eluted fractions were analyzed by SDS-PAGE followed by autoradiography. As shown in Figures 1 and 2, in the incompletely unfolded state in which the protein derived from RuGli cells exists under non-reducing conditions, two protein bands migrating at positions corresponding to the apparent molecular weights of 150 kD and 120 kD were seen.

A more accurate molecular weight determined by SDS-PAGE under reducing conditions indicates the molecular weights of these subunits are about 165 and 140 kD, respectively, because, when reduced, the larger of the subunits gives rise to two polypeptides, having molecular weights of 135 kD and 35 kD. The smaller component appears as a 35/30 kD doublet in this analysis. While not wishing to be bound by this explanation, it is believed that the 30 kD polypeptide is a cleavage fragment of the 35 kD polypeptide.

10 A laminin binding protein with subunits similar to those of the RuGli cell protein was also isolated from human MG-63 osteosarcoma cells and placental tissues. This human laminin receptor is also composed of two polypeptides. However, the larger subunit has a native molecular weight
15 of 170,000 D.

As used herein, the term "laminin adhesion receptor" (also termed "adhesion receptor for laminin") refers to a cell surface receptor which functions in mediating the adhesion of cells to laminin. It selectively binds to
20 laminin or laminin fragments that retain cell attachment promoting activity and does not bind substantially to vitronectin, fibrinogen, collagen type I or albumin. The native laminin adhesion receptor is composed of two subunits having molecular weights of about 165 to 170 kD
25 and about 140 kD. The larger polypeptide in turn is composed of two disulfide linked chains with molecular weights of about 135 kD and 35 kD.

The term "laminin adhesion receptor" refers to both the native structures and modifications or isoforms of those
30 structures which retain laminin or laminin-fragment binding activity. Laminin or laminin-fragments retaining cell attachment promoting activity are referred to as the ligand of the laminin adhesion receptor. It is understood that limited modifications may be made to the structure of the
35 laminin adhesion receptor without destroying the ligand binding activity and that only a portion of the entire

primary structure may be required in order to effect activity. Fragments of the receptor that retain the ligand binding activity are included within the definition.

"Substantially pure," when used to describe the state of 5 the laminin adhesion receptor of the invention, denotes the receptor essentially free of other proteins normally associated with or occurring with the receptor in its native environment and substantially separated from interfering and diluting cell surface proteins or other 10 proteins, such as antibodies, introduced through, for example, isolation procedures. More than one isoform of the receptor may be present, however, in the "substantially pure" preparation.

The heterodimeric structure of the isolated laminin 15 adhesion receptor, the disulfide-bonded two-chain composition of its larger subunit and the increase in molecular weight of the smaller subunit upon reduction are all characteristic of the class of receptors termed integrins (See Ruoslahti and Pierschbacher, Science 238:491 20 (1987), which is incorporated herein by reference). Integrins, which include a number of receptors, share amino acid sequence similarities.

To explore the relationship of the laminin adhesion receptor with known integrins, the receptor was 25 immunoblotted with affinity-purified polyclonal antibodies to the fibronectin and vitronectin receptors. The anti-fibronectin receptor antibody cross-reacted with the smaller (beta) subunit of the laminin adhesion receptor, whereas no reactivity was seen with the anti-vitronectin 30 receptor antibody. These results show that the laminin adhesion receptor is a distinct protein, one subunit of which may be related to the fibronectin receptor beta subunit. The α subunit of the laminin receptors was reactive with an antibody against a peptide modelled after 35 the known sequence of the cytoplasmic domain of the α_3 subunit (Hynes et al., J. Cell Biol., in press), suggesting

that the α subunit of the laminin receptor is α_3 . As shown in Figure 3, the β subunit is closely related to the fibronectin receptor β subunit. However, Figure 4 shows that it can differ from it immunologically, at least in the case of the RuGli cell receptors.

The ligand binding specificity of the laminin receptor was studied in liposome binding assays. The receptor was incorporated into phosphatidylcholine liposome membranes by the method of Pytela et al. Cell 40:191, (1985), which is incorporated herein by reference. Briefly, a detergent solution containing the receptor and a phospholipid are dialyzed against a detergent-free buffer. The resulting liposomes adhered strongly to substrates coated with laminin. They also bound, to a variable extent, to fibronectin and type IV and type I collagen, but not to substrates coated with vitronectin, fibrinogen or albumin. In contrast, liposomes made with fibronectin receptor from the same cells bound strongly to fibronectin and showed no interaction with laminin. The attachment of the liposomes prepared with the laminin adhesion receptor was inhibited by purified cell-attachment promoting laminin fragments, but not by the GRGDSP peptide that inhibits the binding of cells to fibronectin and vitronectin (Gehlsen et al. J. Cell. Biol. 106:925-950 (1988), which is incorporated herein by reference) nor another peptide derived from laminin, YIGSR (Graf et al., Cell 48:989, (1987), which is incorporated herein by reference). All peptides are identified by their conventional one letter abbreviation. These results show that the laminin adhesion receptor can incorporate into the liposome membranes and that it conveys to liposomes the expected binding to laminin.

The laminin adhesion receptor liposomes are useful for targeting liposomes into basement membranes. For instance, deposits in the kidney glomeruli may be treated with liposomes loaded with proteolytic and other enzymes by using receptor-mediated targeting.

To coat a nonlipid surface with receptors, a receptor from a solution is adsorbed or covalently coupled to a surface such as plastic using one of the many well known methods available for such coated or coupling. Receptor fragments that retain the ligand-binding activity but lack the membrane embedded portion of the molecule are advantageously used as they are more soluble than the complete receptor. Such materials coated with receptors are useful as prostheses where attachment of basement membrane is desired, such as, for example, an artificial lens.

The binding specificities of the isolated laminin receptors have also been examined in a radioreceptor assay that consists of incubating ^{125}I labeled receptor in microtiter wells coated with laminin or other proteins and the measurement of the binding of radioactivity to the wells. The laminin receptors from each of the cell types bound to wells coated with laminin. They all showed also some affinity for fibronectin and some of the receptors bound to type IV collagen (Figure 6). The latter result suggests that the receptors isolated from the various cell types fall into two categories - type IV collagen non-binding (for example, RuGli) and type IV collagen binding (for example, MG63). No chemical difference has been established between these types of receptors as yet.

The radioreceptor assay was also used to assess the ability of various protein fragments to inhibit the interaction between laminin and its receptor (Figure 7). Fragments of laminin derived from the portion made up of the COOH-terminal parts of the molecule (Dillner et al., *supra*) inhibited the binding. Moreover, fibronectin fragments derived from the cell attachment region of that molecule were also inhibitory. The ability of the fragments to inhibit the laminin receptor interaction will be useful in preventing the attachment of tumor cells to blood vessel basement membranes. The assay will be useful in identifying other substances capable of inhibiting the

laminin receptor-laminin interaction.

Monoclonal and polyclonal antibodies against the laminin adhesion receptor were prepared according to procedures well known in the art. The polyclonal antibodies are
5 absorbed with the purified fibronectin receptor coupled to Sepharose (Argraves et al., supra). Since the beta subunits of the laminin and fibronectin receptors are similar and since the fibronectin receptor beta subunit is shared by a number of other integrins, this treatment will
10 eliminate antibodies directed against determinants shared by the receptors in this integrin family. It will also eliminate antibodies against contaminating proteins that would bind nonspecifically to both the laminin and fibronectin affinity matrices. If necessary, such
15 antibodies can also be removed by absorbing the antiserum with "a mock-purified receptor preparation" (see Example VI). The specificity of the antibodies is examined by carrying out enzyme immunoassays, immunoprecipitation of detergent extracts from surface-iodinated and metabolically
20 labeled cells and immunoblotting of isolated laminin receptor, other isolated integrins and whole cell extracts by methods well known in the art. As shown in Figure 4, in the case of the RuGli cell receptors, the absorption of an antiserum made against the RuGli cell laminin receptor with
25 the RuGli cell fibronectin receptor yielded an antibody preparation that is primarily reactive with the laminin receptor.

Monoclonal antibodies are prepared by immunizing with the isolated receptor or other material containing the
30 receptor followed by isolation of antibody-producing hybridoma cells, as is well known in the art. (See, for example, Harlow and Lane, ANTIBODIES: A LABORATORY MANUAL, Cold Spring Harbor, 1988, which is incorporated herein by reference.) The appropriate hybridoma cells are selected
35 by performing enzyme immunoassays with the purified laminin receptor. Antibodies specific for the laminin receptor are obtained by using immunoblotting to select

those antibodies that are reactive with either one of the subunits of the laminin receptor but do not react with other integrin subunits. To obtain antibodies specific for the laminin receptor that can inhibit the laminin-binding activity of the receptor, the subunit-specific monoclonal antibodies are tested as inhibitors of cell attachment to laminin. This assay is performed by allowing cells to attach to microtiter wells in the presence and absence of the antibody. Finally, selection of antibodies that promote attachment is performed by coating microtiter wells with purified antibody at various concentrations followed by addition of cells possessing the laminin receptor into the wells. Increased attachment of the cells relative to wells coated with an inert protein such as serum albumin shows the presence of antibody with the desired properties.

The laminin adhesion receptor can also be used to screen various compounds for their ability to compete for binding to laminin. Preferably, such screening is accomplished using an inhibition assay, such as that described in Example VIII, although other methods will be obvious to one skilled in the art. The inhibition assay can be used advantageously to screen unknown substances or mixtures of substances for inhibitory activity. If such a mixture is found to have inhibitory activity, it can then be fractionated in order to isolate or identify the particular active compound. Such active compounds can then be used either to promote attachment to a substrate of cells exhibiting laminin adhesion receptors by coating the substrate with such a compound or to inhibit adhesion of such cells to laminin by providing the compound in soluble form.

The following examples are intended to illustrate but not limit the invention. While they are typical of those that might be used, other procedures known to those skilled in the art may be alternatively employed.

EXAMPLE I

ISOLATION OF LAMININ RECEPTORa. RuGli Cells

5 Glioblastoma cell line RuGli, obtained from the Max Planck Institute, Tubigen, West Germany, was used as a source of rat laminin adhesion receptor. Cells were grown to confluency and the cell surfaces were iodinated with ^{125}I by the lactoperoxidase method (Lebien et al, J. Immunol. 129:2287 (1982), which is incorporated herein by
10 reference) and extracted with Tris-buffered saline, pH 7.2, containing 25 mM octyl-beta-D-thioglucoside (Calbiochem, La Jolla, CA), 1 mM MnCl_2 , and 1 mM phenylmethylsulfonylfluoride (PMSF) (extraction buffer).
15 Mn^{2+} was included in the buffer because it facilitates the isolation of the fibronectin receptor by fibronectin affinity chromatography.

Laminin fragments having cell attachment promoting and neurite promoting activity were isolated from human
20 placenta by the immunoaffinity chromatography method of Wewer et al., J. Biol. Chem. 258:12654 (1983), which is incorporated herein by reference. The 4E10 anti-human laminin antibody that was used for isolation is commercially available from Telios Pharmaceuticals, Inc.,
25 San Diego, CA.

A laminin-Sepharose column was prepared by coupling 90 mg of human laminin fragments, obtained as above, to 10 ml of cyanogen bromide-activated Sepharose gel (Pharmacia, Uppsala, Sweden) following the manufacturer's instructions.

30 The soluble extract from 5 ml of packed surface-iodinated cells was applied to a 10 ml (bed volume) laminin-Sepharose column. The cell extract was passed through the column two times and the column was washed with 30 ml of the extraction buffer. An initial dilution was
35 done with 20 ml buffer containing 2 mg/ml of the synthetic

peptide GRGDSP followed by 20 ml of cation-free buffer containing 20 mM EDTA. Aliquots of each collected 2 ml fraction were electrophoresed on a 7.5% SDS polyacrylamide gel under non-reducing conditions. Protein bands were visualized by overnight autoradiography. The GRGDSP elution released no specifically eluted bands from the affinity matrix, whereas the EDTA elution released a laminin adhesion receptor as shown in Figure 1.

As shown in Figures 1 and 2, in the incompletely unfolded state in which the protein derived from RuGli cells exists under non-reducing conditions, two protein bands migrating at positions corresponding to the apparent molecular weights of 150 kD and 120 kD were seen. A more accurate molecular weight determined by SDS-PAGE under reducing conditions indicates the molecular weights of these subunits are about 165 and 140 kD, respectively, because, when reduced, the larger of the subunits gives rise to two polypeptides, 135 kD band and a 35/30 kD doublet. While not wishing to be bound by this explanation, it is believed that the 30 kD polypeptide is a cleavage fragment of the 35 kD polypeptide.

b. MG-63, A-431, and U-251 Cells

Human osteosarcoma cells (MG-63; ATCC Accession No. CRL 1427) human epidermal carcinoma cells (A-431; ATCC Accession No. CRL-1555) and human glioblastoma cells (U-251) were used as sources of human integrin-type laminin receptors. The receptors were isolated by the methods described above for the RuGli receptor. The receptor obtained had the same electrophoretic appearance as the RuGli receptor shown in Figure 1, except that the larger (α) subunit migrated somewhat more slowly than the RuGli receptor α subunit giving an apparent molecular weight of about 170,000D.

EXAMPLE II

INCORPORATION OF LAMININ RECEPTOR INTO LIPOSOMES
AND BINDING OF THE LIPOSOMES TO SUBSTRATES

Phosphatidylcholine liposomes incorporating the cell
5 surface receptors were prepared essentially by the method
of Mimms et al., Biochemistry 20:883, (1981), as described
for the fibronectin receptor (Pytela et al., Meth. Enzymol.
144:475-489, 1987), both references being incorporated
herein by reference. Egg yolk phosphatidylcholine (Sigma,
10 St. Louis, MO) and ³H-phosphatidylcholine (New England
Nuclear, Boston, MA), were dissolved in the receptor
fractions at 100 µg/ml and this solution was then dialyzed
against 50 mM Tris-HCl containing 500mM NaCl, 1 mM Ca₂Cl
and 1 mM Mg₂Cl for 24 hours at 4°C. The resulting
15 liposomes were isolated by floating to the surface of a
sucrose gradient in ultracentrifugation and tested for
binding of various substrates.

The substrates were prepared by coating polystyrene
microtiter plate wells (Linbro/Titertek, Inglewood, CA)
20 with extracellular matrix proteins including laminin,
fibronectin, collagens and vitronectin (Telios
Pharmaceuticals, Inc., La Jolla, CA), according to the
method of Engvall et al., J. Cell. Biol. 103:2457 (1986),
which is incorporated herein by referenc, using 20 µg/mL in
25 PBS overnight at 4°C. Coating with bovine serum albumin
(BSA; 5 mg/mL) was used to determine nonspecific background
binding (Sigma Chemical Co., St. Louis, MO). The laminin
receptor-liposome preparations showed strong dose-dependent
binding to microtiter wells coated with laminin, bound
30 slightly to fibronectin, and showed no binding over the
background to collagens type I or IV or to vitronectin, as
shown in Figure 5.

The fibronectin receptor isolated from the same RuGli
cell extract was used for comparison in the liposome assay.
35 Liposomes prepared with this receptor bound to a greater
degree to fibronectin than the laminin receptor liposomes

and showed no binding to laminin. Further specificity controls included inhibition of the laminin receptor binding by adding cell attachment-promoting chymotryptic human laminin fragments to the liposome assay. These 5 fragments inhibited the binding of the laminin receptor liposomes to laminin, whereas the GRGDSP peptide or the YIGSR laminin peptide did not. Representative results from these assays are shown in Figure 5.

EXAMPLE III

10 USE OF THE RECEPTOR-LIPOSOME PREPARATION TO TARGET THERAPEUTIC AGENTS

Receptor liposomes are prepared as described above and a desired therapeutic agent is incorporated into them according to published methods (Gregoriadis and Senior, 15 Biochem. Soc. Trans. 12:337, (1984) which is incorporated herein by reference). These conjugated liposomes are injected intravenously using 1-300 μ mol of lipid per kg of body weight or applied locally. Such laminin receptor liposomes are used for example, to target drugs to tissues 20 that contain large amounts of laminin.

EXAMPLE IV

ANALYSIS OF RECEPTOR SPECIFICITY BY RADIORECEPTOR ASSAY

The binding of laminin receptor to laminin and the 25 effect of antibodies and protein fragments on this binding was examined in a radio-receptor assay previously described for the fibronectin receptor (Hautanen, et al., J. Biol. Chem. 264:1437-1442 (1989)). Microtiter wells were coated with 1-2 μ g/ml of human laminin, bovine type I collagen 30 (Collaborative Research, Lexington, MA), mouse type IV collagen (BRL, Bethesda, MD), human plasma fibronectin or bovine serum albumin. 125 I-labeled receptor isolated from surface iodinated cells was allowed to bind to the coated wells for 2 hours at room temperature in the presence of

Tris-buffered saline (150mM NaCl, 50mM Tris-HCl, pH 7.5, TBS) containing 50mM octyl- β -glucopyranoside, 1mM PMSF, and 1mM MnCl₂. The amount of receptor added per well was equivalent to 10⁴ cpm. Following incubation, wells were washed, and bound receptor was solubilized with 1% sodium dodecyl sulfate (SDS) in TBS, and quantitated by counting the bound radioactivity. Nonspecific binding was measured in wells coated with albumin. The total binding varied between 5 and 10% of the added radioactivity, and 70-80% of this was specific by the above criteria. The laminin receptors from each of the cell types tested bound to laminin in the assay as shown in Figure 6. The receptors also bound to varying degrees to fibronectin and type IV collagen. The reasons for this variation are not known. Without wishing to be bound by that explanation, it is believed that the difference may be in the β subunit of the receptor such that the receptor with affinity for type IV collagen has the composition $\alpha_3\beta_1$, (See Ruoslahti and Pierschbacher, supra) and the one with no affinity for type IV collagen has a different β subunit. Support for this position is provided by the immunological differences between the RuGli cell laminin and fibronectin receptors documented in Figure 4.

EXAMPLE V

25 LOCALIZATION OF THE LAMININ RECEPTOR BINDING SITE

a. Electron Microscopy

Electron microscopic analysis of laminin receptor-laminin complexes was performed as described for receptor-fibronectin complexes (Gallit and Ruoslahti, J. Biol. Chem. 263:12927 (1988), which is incorporated herein by reference) with some modifications. Receptor preparations from affinity chromatography eluates were mixed with either mouse laminin (BRL, Bethesda, Md) or with pepsin fragments of human laminin (Dillner, et al., Exp. Cell. Res. 177:186 (1988), which is incorporated herein by reference) in the

presence of 1 mM each of Ca^{2+} , Mg^{2+} and Mn^{2+} . Various ratios of receptor and laminin were used, and all incubations of the receptor-laminin complexes were performed at 4°C to avoid the aggregation of laminin known to take place in the presence of divalent cations at higher temperatures.

When the laminin receptors were mixed with mouse laminin, some of the laminin molecules appeared to have complexed with the receptor. The receptor was bound to the globular end of the laminin long arm in these complexes. No structures resembling the laminin receptor complexes were observed in control preparations consisting of laminin alone or laminin mixed with the MG-63 or RuGli cell fibronectin receptor. Examination of a large number of fields from several independently prepared samples showed that among more than 100 possible receptor-laminin complexes, approximately 80% had the receptor associated with the globular domain at the end of the long arm. Among the remainder of the possible complexes, the receptor molecule appeared to be associated variously with one of the short arms or the center of the cross. Laminin-receptor complexes similar to those obtained with mouse laminin were seen when pepsin fragments of human laminin were mixed with the laminin receptor.

25 b. Localization with Monoclonal Antibodies

The preferential binding of the receptor to the globular domain at the end of the long arm of the laminin molecule was similar to what has previously been shown for a group of monoclonal antibodies capable of inhibiting the neurite-promoting activity of laminin. Monoclonal antibodies against pepsin-extracted human laminin were prepared by the method of Engvall et al., J. Cell Biol. 103:2457-2465 (1986), which is incorporated herein by reference. The ability of two of these antibodies, 3E5 and 4E10, to interfere with the binding of the receptor to laminin in a radio-receptor assay was examined. The 3E5 and 4E10

antibodies inhibited the binding of the RuGli receptor to laminin in a concentration-dependent fashion. Antibody 2E8, which has no effect on neurite outgrowth and which binds to a region near the center of the laminin cross, 5 showed little inhibition at comparable antibody concentrations. A monoclonal antibody of unrelated specificity was also ineffective. The binding of the MG-63 cell receptor was similarly inhibited by the 3E5 and 4E10 antibodies in this assay, indicating that both laminin 10 receptors may bind to the same site.

c. Localization of monoclonal antibody binding site by cDNA cloning

Laminin cDNA clones were isolated to test proteins produced by these clones for the binding of the monoclonal 15 antibodies that defined the receptor binding site. Screening of placental and endothelial cell λ gt11 cDNA libraries with polyclonal and monoclonal antibodies to laminin yielded numerous reactive clones. DNA sequencing and comparison to the previously published laminin 20 sequences (Pikkarainen, et al., J. Biol. Chem. 262:10454-10462 (1987) and Pikkarainen, et al., J. Biol. Chem. 263:6751-6758 (1988)) showed that several of these clones coded for COOH-terminal portions of the B1 and B2 chains. No A chain clones were identified.

25 Fusion proteins expressed from the B1 and B2 chain clones were tested for reactivity with the monoclonal antibodies in macroplaque filter assay (Suzuki, et al., EMBO J. 4:2519-2524 (1985)). The fusion proteins from all clones encoding the COOH-terminus of the B1 chain reacted 30 with the 3E5 antibody, whereas this antibody did not react with the B2 chain fusion proteins. The shortest clone reactive with the 3E5 antibody encodes the 252 COOH-terminal amino acids of the B1 chain. None of the fusion proteins from the COOH-termini of the B1 and B2 chains, 35 were reactive with the other antibody 4E10.

EXAMPLE VI

PREPARATION OF ANTIBODIES TO LAMININ RECEPTOR
AND THEIR USE TO DETECT THE RECEPTOR

a. Polyclonal antibodies

5 Rabbits were immunized with purified laminin receptor in Freund's complete adjuvant. Booster injections containing 50 μ g of receptor in Freund's incomplete adjuvant were given at three week intervals and serum is collected one week after the last injection. The antiserum was absorbed
10 with human plasma-proteins, fibronectin and laminin, each coupled to Sepharose. In addition, absorption with a "mock-purified receptor" that was obtained by fractionating the same cell or tissue extract that served as the source of the laminin receptor on a Sepharose column to which
15 bovine serum albumin was coupled. The same procedure as in the receptor isolation was followed and those fractions that in the receptor isolation contained the receptor were collected as the mock purified receptor. The antiserum was then further absorbed with fibronectin receptor, also
20 coupled to Sepharose. The fibronectin receptor was isolated by the method of Pytela et al., Supra. The absorption with the mock purified receptor-Sepharose and fibronectin receptor-Sepharose was carried out by passing 2 ml of serum sequentially through columns of 1 ml Sepharose
25 containing 100 μ g of the mock-purified receptor proteins or 100 μ g of the fibronectin receptor. The nonbound material was tested for its reactivity by solid phase enzyme² immunoassay (ELISA; Engvall, Meth. Enzymol. 70:419-439, 1980 which is incorporated herein by reference) with the
30 purified laminin receptor and fibronectin receptor as antigens, and by SDS-PAGE immunoblotting against the receptors and against a whole cell extract or tissue extract from the cells or tissue that served as the source of the receptor. The absorptions were repeated until these
35 assays showed that the antiserum is specific. The antiserum was considered specific for the laminin receptor when it reacted only with the laminin receptor in ELISA,

and bound only to the laminin receptor subunits but not to the fibronectin receptor subunits in immunoblotting, performed according to the method of Towbin et al., Proc. Natl. Acad. Sci. USA 76:4350 (1979) and Argraves et al, 5 supra, (see Figure 4), which is incorporated herein by reference. The same procedure was used to obtain fibronectin receptor specific antibodies from the fibronectin receptor antiserum by absorption with the mock-purified receptor proteins and the laminin receptor. The 10 receptor-Sepharose columns were regenerated by washing with 8 M urea in 50 mM Tris-HCl, pH 7.0.

b. Monoclonal antibodies

Mice were immunized with 10 μ g of purified laminin receptor per injection with the first injection given 15 subcutaneously in Freund's complete adjuvant and the booster injection intraperitoneally without adjuvant. Spleen cells from the immunized mice were collected 3 days after the booster injection and used to generate hybridomas as described in the literature. Hybridomas secreting 20 antibody that reacts with the laminin receptor were selected by solid phase enzyme immunoassay with the laminin receptor as the antigen (Engvall, et al., Meth. Enzymol. 70:419-439, 1980). For the assay, microtiter wells were coated with 1 μ g/ml of receptor solution diluted from the 25 octylglucoside-containing stock solution at least 1:10 to phosphate-buffered saline. Those antibodies that were specifically reactive with the laminin receptor were further selected by testing the positive antibodies against the fibronectin receptor. This reactivity was verified for 30 those antibodies that were reactive with the SDS-denatured receptor by SDS-PAGE immunoblotting against cell extracts containing receptors other than the laminin receptor.

Further selection of antibodies that inhibit the laminin-binding function of the receptor was carried out in 35 cell attachment assays. The monoclonal antibodies were isolated from the hybridoma culture media by affinity

chromatography on protein A-Sepharose (Pharmacia, Uppsala, Sweden), although other methods well known in the art can be used and the isolated antibody was tested for its ability to inhibit the attachment of cells that served as the source of the immunizing receptor to laminin. Microtiter wells were coated with 1 μ g/ml of cell attachment-promoting human laminin fragments and the RuGli cells were added to the wells with various concentrations of each monoclonal antibody. Polyclonal anti-laminin receptor antibodies were used as a positive control and anti-fibronectin receptor alpha subunit monoclonal antibodies as a negative control. Fewer cells were attached to the wells at the end of the assay when the antibody had the desired inhibitory activity.

15

EXAMPLE VII

TESTING OF ANTI-LAMININ RECEPTOR ANTIBODIES IN
TUMOR INVASION ASSAYS

To test for the ability of anti-laminin alpha subunit antibodies to interfere with the invasion of tumor cells through tissue, an amniotic membrane invasion assay is used. RuGli glioblastoma cells are allowed to migrate through an amniotic membrane as described in the literature (Gehlsen et al., J. Cell Biol. 106:925-930, 1988 which is incorporated herein by reference) in the presence of various concentrations of the anti-laminin receptor antibodies and the same control antibodies described above. Fewer cells will migrate through the amniotic membrane in the presence of inhibitory antibodies than in the controls.

30

EXAMPLE VIII

INHIBITION OF LAMININ RECEPTOR
IN THE RADIO-RECEPTOR BINDING ASSAY

Microtiter wells were coated with laminin at 1 to 2 μ L in PBS as described in Example IV. Laminin receptor

isolated from A-431 or RuGli cells as described in Example I was then added to the microtiter wells either alone (LM) or in the presence of chymotrypsin fragments of laminin (chymo. frag., 25 μ g/mL Dillner, et al., Supra), the 110 kD fragment of fibronectin (110 kD frag., 25 μ g/mL, 120 kD fragment in Pierschbacher et al., Cell 26:259-267 (1981)) or unlabelled laminin receptor (Cold LMR).

The results are shown in Figure 7. Data represent the percentage of laminin receptor bound compared to uninhibited laminin binding at 100 percent. As can be seen, cell attachment promoting fragments of laminin derived from the portion made up of the COOH-terminal parts of the molecule inhibited the binding of the laminin receptor to the laminin coated wells. Moreover, the fibronectin fragment, which is derived from the cell attachment region of that molecule, was also inhibitory. As expected, unlabeled laminin receptor, which was tested as a control, also inhibited the binding of the radioactive receptor.

20

Industrial Application

This invention finds direct and immediate application in the assay of laminin receptor in cells and tissues. The isolation method described above can be used to assay cultured cells or tissue samples for their content of the laminin receptor. Such analysis will be important in determining the adhesion capacity of cells such as those in tumors. Alternatively, the isolated receptor can be used to prepare antibodies for receptor quantitation. Together with such antibodies, the receptor will permit establishment of assays for the receptor such as a radioimmunoassay or ELISA. Compounds binding to the receptor so that they compete with the binding of the receptor to laminin can be selected with the receptor and are useful, for example, in preventing attachment of the tumor cells to basement membranes during tumor invasion. Alternatively, such substances, when presented in an

ininsoluble form may be used to reproduce the cell attachment promoting effect of laminin in tissue reconstruction. A reagent consisting essentially of the laminin receptor described in the foregoing examples may be
5 used to carry substances to selected tissues.

Although the invention has been described with reference to the presently-preferred embodiments, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly,
10 the invention is limited only by the following claims.

WE CLAIM:

1. A substantially pure active laminin adhesion receptor, characterized by comprising two subunits, one of which is α_3 , the α_3 subunit further comprising two disulfide linked polypeptides, said receptor selectively
5 binding with laminin or cell attachment-promoting fragments of laminin.

2. The receptor of claim 1, wherein said mammalian is a rat.

10 3. The receptor of claim 1, wherein said mammalian is a human.

4. The receptor of claim 1 wherein said receptor can be eluted from said selective binding with laminin or cell attachment-promoting fragments of laminin with ethylene-diaminotetraacetic acid.

5. A composition of matter comprising the receptor of claim 1 incorporated into lipid vesicles.

6. A composition of matter comprising the receptor of claim 1 coated onto a substrate.

7. A composition of matter comprising fragments of the receptor of claim 1 which retain ligand binding activity.

8. A method for isolating a laminin cell surface receptor from a mammalian cell extract preparation, comprising the steps of:

- 5
- a. providing the cell attachment-promoting portion of laminin;
 - b. coupling said material to a matrix to constitute an affinity column;
 - c. running said cell extract preparation through the column in the presence of divalent cation

so as to achieve binding between the cell surface receptors to be isolated and laminin cell attachment site;

- 5 d. eluting said column with a solution containing a substance capable of selectively eluting said cell surface receptor; and
- e. collecting said cell surface receptors eluted thereby.

9. Receptors isolated by the method of claim 6.

10. The method of claim 8, wherein the affinity matrix is Sepharose^R.

11. The method of claim 8, wherein said substance comprises the cell attachment site of laminin defined by the attachment of RuGli glioblastoma cells and MG-63 osteosarcoma cells.

12. The method of claim 8, wherein said cell extract preparation is derived from cultured mammalian cells.

13. The method of claim 8, wherein said cell extract preparation is derived from animal tissue.

14. The method of claim 8, wherein said cell extract preparation is derived from human cells.

15. The method of claim 14, wherein said human cells are osteosarcoma cells.

16. The method of claim 8, wherein said cell extract preparation is derived from placental tissue.

17. The method of claim 8, wherein said cell extract preparation is derived from rat cells.

18. Polyclonal antibodies against the laminin adhesion receptor of claim 1.

19. The antibodies of claim 18 wherein said antibodies are reactive with the alpha subunit of said laminin adhesion receptor.

20. Monoclonal antibodies made by immunizing with the purified laminin receptor of claim 1.

21. The antibodies of claim 20 wherein said antibodies are reactive with the alpha subunit of said laminin adhesion receptor.

22. Antibodies reactive with the β chain of the laminin adhesion receptor but not reactive with β_1 .

23. A method of determining the presence of the laminin adhesion receptor in a sample, comprising the steps of:

- a. providing antibodies reactive with the laminin adhesion receptor;
- 5 b. contacting said sample with said antibodies; and
- c. determining binding between said antibodies and components of said sample, wherein said binding indicates the presence of said laminin
10 receptor in said sample.

24. The method of claim 23 wherein said antibodies are polyclonal antibodies.

25. The method of claim 23 wherein said antibodies are monoclonal antibodies.

26. A method of targeting agents to specific tissue components, comprising:

- a. incorporating laminin adhesion receptor into liposomes having agents contained therein;
- 5 b. presenting said incorporated liposomes to said

specific tissue components.

27. A method of screening for compounds reactive with the laminin adhesion receptor comprising the steps of:

- a. contacting ligands with said receptor; and
- b. determining the reactivity of said compounds with said receptor.

5

28. A method of inhibiting the binding of the laminin adhesion receptor to laminin by providing compounds reactive with the laminin adhesion receptor.

29. A method of inhibiting the attachment to a substrate of cells exhibiting laminin adhesion receptors, comprising providing solubilized compounds reactive with the laminin adhesion receptor.

30. A method of promoting the attachment to a substrate of cells exhibiting laminin adhesion receptors, comprising providing insoluble compounds reactive with the laminin adhesion receptor.

1/4

FIG. 1





FIG. 2



2 / 4




FIG. 4




FIG. 3

3 / 4




FIG. 5

4 / 4




FIG. 6




FIG. 7

INTERNATIONAL SEARCH REPORT PCT/US89/02240

International Application No.

I. CLASSIFICATION OF SUBJECT MATTER		
According to International Patent Classification (IPC) or to both National Classification and IPC INT. CL. 4 A61K 9/64; C07K 3/20, 15/14, 17/00 17/02; GOIN 33/567, 574, 577 U.S.C1. 424/450; 436/503, 547, 548, 813; 530/387, 395, 413, 810, 812, 828		
1. FIELDS SEARCHED		
Minimum Documentation Searched?		
Classification System	Classification Symbols	
INT C1.	A61K 9/64	
US. C1.	424/450; 436/503, 547, 548, 813; 530/387, 395, 413, 810, 812 828.	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched		
APS: Laminin and (receptor? or binding (w)Protein?) Biosis: Laminin(2A)(receptor? or binding(w)protein?)) and (antibod? or antisera? or affinity or purif? or isolat?)		
III. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X, P	Science, vol. 241, no.10, issued 2 September 1988. K,R. GEMLSEN ET AL, "The human laminin receptor is a member of the integrin family of cell adhesion receptor" pages 1228-1229. See entire document.	1-6, 8-17, 22
Y	METHODS IN ENZYMOLOGY, vol. 144, issued 1987, R. PYTELA ET AL, "Arginine-glycine- aspartic acid adhesion receptors", pages 475-489. See pages 479-483 and 489.	1-30
* Special categories of cited documents: ¹⁰ "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such docu- ments, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
01 August 1988	17 OCT 1989	
International Searching Authority	Signature of Authorized Officer	
ISA/US	David A. Saunders	

III DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)			
Category	Citation of Document	With indication, where appropriate, of the relevant passages	Relevant to Claim No.
X Y	Cell, vol. 48, no. 6, issued 27 March 1987, J. GRAF ET AL, "Identification of an amino acid sequence in laminin mediating cell attachment, chemotaxis, and receptor binding", pages 989-996. See page 990 cols. 1 and 2, and page 992, col. 2		27-30 1-17
Y	Biochemical Society Transactions, vol. 12, issued 1984, G. GREGORIADIS ET AL, "Targeting of small unilamellar liposomes to the galactose receptor in vivo", pages 337-339. See page 337, col. 1.		26
X Y	Biological Abstracts, vol. 80, No. 5 issued 1 September 1985, P. MAND ET AL, "Expression of laminin receptor in normal and carcinomatous human tissues as defined by a monoclonal antibody", page 587, col. 1, the abstract no. 41870, Cancer Research, 1985, 45(6), pages 2713-2719. See entire abstract.		23, 25 20-22
Y	The Journal of Biological Chemistry, vol. 258, no. 20, issued 25 October 1983, U. WEWER ET AL, Human laminin isolated in a nearly intact biologically active form from placenta by limited proteolysis", pages 12654-12660. See abstract and page 12659, col. 2.		1-17
X Y	US, A, 4,565,789 (LIOTTA ET AL) 21 January 1986. See col. 5, line 53-col. 6, line 4; col 6, lines 1-27 and 48-55; col 10, lines 40-63; col. 15, lines 1-12.		23-25, 27-29 1-22
X Y	Proceedings of the National Academy of Sciences USA, vol. 29 no. 18, issued September 1987, NR. SMALHEISER ET AL, "Cranin: A laminin-binding protein of cell membranes", pages 6457-6461. See Abstract and page 6461, col. 2.		1, 7, 18, 19 23, 24 2-6, 8-17, 20-22-25-30
X Y	Biological Abstracts, vol. 85, no. 2, issued 15 January 1988, U.M. WEWER ET AL, "Role of laminin receptor in tumor cell migration", the abstract no. 17879, Cancer Research, 4987, 47(21) pages 5691-5698. See entire abstract.		23, 24 18, 19

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE ¹

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____ because they relate to subject matter ¹² not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out ¹³, specifically:

3. ☐ Claim numbers _____, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING ²

This International Searching Authority found multiple inventions in this international application as follows:

See attachment

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application. Telephone practice

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
☐ No protest accompanied the payment of additional search fees.

ATTACH TO FORM 210, SUPPLEMENTAL SHEET (2), PART VI

Group I, claims 1-17, drawn to a receptor protein and its method of isolation.

Group II, claims 18-25, drawn to antibodies to the receptor and uses thereof.

Group III, claim 26, drawn to targetting methods using liposomes bearing the receptor.

Group IV, claims 27-30, drawn to binding methods using the receptor.

The inventions listed as groups II-IV do not meet the requirements for Unity of Invention for the following reasons: Groups II-IV represent alternate methods of using the product of Group I. That is the product of Group I is used to make antibodies, to make targettable liposomes, or as a receptor in binding studies. Unity of Invention requires only one use.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record.**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.